

## **Informed Consent for Cataract Surgery**

### **WHAT IS A CATARACT AND HOW IS IT TREATED?**

The lens in the eye can become cloudy and hard, a condition known as a cataract. Cataracts can develop from normal aging, from an eye injury, or if you have taken medications known as steroids. Cataracts may cause blurred vision, dulled vision, sensitivity to light and glare, and/or ghost images. If the cataract changes vision so much that it interferes with your daily life, the cataract may need to be removed. Surgery is the only way to remove a cataract. You can decide not to have the cataract removed. If you don't have the surgery, your vision loss from the cataract will continue to get worse.

### **HOW WILL REMOVING THE CATARACT AFFECT MY VISION?**

The goal of cataract surgery is to correct the decreased vision that was caused by the cataract. During the surgery, the ophthalmologist (eye surgeon) removes the cataract and puts in a new artificial lens called an intraocular lens or IOL. Cataract surgery will not correct other causes of decreased vision, such as glaucoma, diabetes, or age-related macular degeneration. Most people still need to wear glasses or contact lens after cataract surgery for either near and/or distance vision and astigmatism.

### **WHAT TYPES OF IOLs ARE AVAILABLE?**

Your ophthalmologist will help you decide on the type of IOL that will replace your cloudy lens. There are IOLs available to treat nearsightedness (myopia), farsightedness (hyperopia), and astigmatism. IOLs usually provide either near or distance vision: these single focus lenses are called monofocal IOLs. Some newer IOLs can provide for near, intermediate, and distance vision: these multiple focus lenses are called multifocal IOLs. IOLs that treat astigmatism are called toric IOLs. You can also have one eye corrected for near vision, and the other for distance vision, a choice called monovision.

### **WHAT IS ASTIGMATISM? ARE THERE OTHER TREATMENTS FOR IT?**

Patients with nearsightedness and farsightedness often also have astigmatism. An astigmatism is caused by an irregularly shaped cornea; instead of being round like a basketball, the cornea is shaped like a football. This can make your vision blurry. In addition to toric IOLs, astigmatism can be reduced by glasses, contact lenses, and refractive surgery (LASIK or PRK). There is also a procedure called a limbal relaxing incision (LRI), which can be done at the same time as the cataract operation, or as a separate procedure. A limbal relaxing incision (LRI) is a small cut or incision the ophthalmologist makes into your cornea to make its shape rounder. Any attempt at astigmatism reduction could result in over- or under-correction, in which case glasses, contact lenses, or another procedure may be needed.

